

## Precipitation of Acute Psychotic Episodes by Intensive Meditation in Individuals with a History of Schizophrenia

BY ROGER WALSH, M.D., PH.D., M.R.A.N.Z.P., AND LORIN ROCHE

The term "meditation" refers to a family of practices that induce the conscious training and directing of attention, with the ultimate aim of controlling the mind and enhancing psychobiological health.

Despite its history as perhaps the oldest psychotherapeutic practice, meditation has gained widespread popularity in the West only within the last two decades and began to receive serious research attention even more recently. However, a rapidly expanding literature on meditation has demonstrated a variety of psychological, physiological, electroencephalographic, and chemical effects, and there now seems little question that meditation is capable of producing a wide range of psychotherapeutic benefits (1).

There have been several reports of adverse meditation effects, including depersonalization, altered real-

ity testing, and the appearance of previously repressed, highly charged memories and conflicts (1-5). However, as yet no clear picture has emerged of the individuals at risk for pathologic reactions.

We have observed several thousand people given intensive meditation training, and we are familiar with three psychotic episodes. We are reporting these cases because they show possible etiologic similarities that may be useful in identifying a population at risk.

### Case Reports

*Case 1.* Mr. A, a 25-year-old man, attended a 1-week meditation retreat that involved many hours each day of sitting and walking meditation and total silence, without communication of any kind (even eye contact). On the 6th day his roommate reported that Mr. A appeared agitated. On the 7th day Mr. A suddenly began to weep and scream; he required physical restraint to avoid hurting himself or others and was clearly psychotic. He was given 150 mg of oral thioridazine and within 4 hours was calm and rational. He then reported that he had been hospitalized four times in the preceding 5 years for acute schizophrenic episodes, that he had been receiving outpatient psychotherapy sporadically, and that he

Received March 12, 1979; accepted April 20, 1979.

Dr. Walsh is Assistant Professor, Department of Psychiatry and Human Behavior, Medical Science 1, University of California, Irvine, Calif. 92717, where Mr. Roche is a graduate student, School of Social Sciences.

had been prescribed a small maintenance dose of phenothiazine, which he had stopped taking. In addition, from the 3rd day of the retreat he had fasted, intensified his meditation as much as possible, and curtailed his sleep to only 1 or 2 hours a night at most. He was returned to his home and did well on small maintenance doses of phenothiazines.

**Case 2.** Ms. B was a 23-year-old woman who attended a 2-week intensive retreat that involved 18 hours of meditation a day. On the 10th day she began asking bizarre questions and accused people of stealing her possessions. When she was interviewed it became apparent that she was undergoing an acute schizophrenic episode. A unique feature was the remarkable lucidity and detachment with which she was able to observe and describe the process she was going through; this was apparently a result of her meditation training. She reported that she had been hospitalized several times in the past for acute schizophrenic episodes and that she had eaten and slept only sporadically over the previous several days. She was given thioridazine and was instructed to stop meditating and to assist the staff in various tasks. Ms. B resumed eating and sleeping, and after 4 days her condition had improved sufficiently so that she was able to return home.

**Case 3.** Ms. C, a 23-year-old woman attending an intensive meditation retreat, began to exhibit bizarre behavior on the 4th day of the retreat. When she was interviewed, Ms. C exhibited suicidal ideation and features of an incipient acute schizophrenic episode, and she reported a history of multiple hospitalizations for schizophrenia. Ms. C was instructed to stop meditating and was given oral chlorpromazine. However, within hours of this interview she made a serious suicide attempt with a knife and was hospitalized involuntarily.

### Discussion

These cases suggest that the combination of intensive meditation, fasting, sleep deprivation, a history of schizophrenia, and the discontinuation of maintenance doses of phenothiazines can be hazardous. Schizophrenic episodes can be precipitated by almost any severe stress, including a range of intensive therapeutic modalities, and the combination of intensive meditation practice, together with sleep and food deprivation, is certainly demanding. On the other hand, some data suggest that meditation in moderation can be useful in treating a range of psychopathology, including schizophrenia (1, 6-8). The population at risk for the syndrome we have described seems to include only people with a history of schizophrenia, for whom maintenance medication, adequate food and sleep, and a less intensive approach to meditation are indicated. However, even psychologically healthy individuals may experience changes in their perception of reality during the initial phases of intensive meditation (1, 9). Such changes are not necessarily pathologic and may

reflect in part a heightened sensitivity to the (usually subliminal) perceptual distortions to which we are all subject (9).

We were interested to note two unique features associated with these episodes. The first was the degree of insight and detachment that Ms. B was able to obtain, especially during interviews, which appeared to be a result of her meditation training. The second was the extraordinarily high degree of skill and empathy with which the retreat staff, all advanced meditators, handled these clients. This seemed to reflect the staff's psychological sophistication and familiarity with altered states. In addition, it is interesting that all three patients recovered rapidly.

An increasing number of people practice brief meditation sessions daily, and tens of thousands of Americans attend intensive meditation retreats each year.<sup>1</sup> Although some of the meditation teaching organizations are beginning to screen retreat applicants for previous psychiatric history, episodes such as those described above are bound to occur from time to time, and clinicians can help by being aware of the population at risk. However, psychotic episodes are rare, and our impression, which is borne out by the classical and research literature (1, 6-8), is that most people show significant gains in a number of areas of psychological well-being.

### REFERENCES

1. Shapiro D, Walsh R (eds): *Meditation: Self Control Strategy and Altered States of Consciousness*. New York, Aldine Press (in press)
2. Otis L: Adverse effects of transcendental meditation, in *Meditation: Self Control Strategy and Altered States of Consciousness*. Edited by Shapiro D, Walsh R. New York, Aldine Press (in press)
3. French AP, Schmid AC, Ingalls E: Transcendental meditation, altered reality testing, and behavioral change. *J Nerv Ment Dis* 161:55-58, 1975
4. Lazarus AA: Psychiatric problems precipitated by transcendental meditation. *Psychol Rep* 10:39-74, 1976
5. Kennedy R: Self-induced depersonalization syndrome. *Am J Psychiatry* 133:1326-1328, 1976
6. Walsh R, Shapiro D (eds): *Beyond Health and Normality: Explorations of Extreme Psychological Wellbeing*. New York, Van Nostrand Reinhold (in press)
7. Glueck BC, Stroebel CF: Biofeedback and meditation in the treatment of psychiatric illness. *Compr Psychiatry* 16:303-321, 1975
8. Shafii M: Adaptive and therapeutic aspect of meditation. *Int J Psychoanal Psychother* 2:367-382, 1973
9. Walsh R: Initial meditation experiences: I. *Journal of Transpersonal Psychology* 9:151-192, 1977

<sup>1</sup>Gallup poll release, Nov. 18, 1976, entitled "Six Million Involved in Transcendental Meditation, Americans Taking up Religious, Spiritual Experimentation."